arXiv:1511.09321v1 [math.CO] 30 Nov 2015

On connected degree sequences

Jonathan McLaughlin

Department of Mathematics, St. Patrick's College, Dublin City University, Dublin 9, Ireland

Abstract

This note gives necessary and sufficient conditions for a sequence of non-negative integers to be the degree sequence of a connected simple graph. This result is implicit in a paper of Hakimi. A new alternative characterisation of these necessary and sufficient conditions is also given.

Keywords: connected graph, degree sequence
2010 MSC: 05C40

1. Introduction

A finite sequence of non-negative integers is called a graphic sequence if it is the degree sequence of some finite simple graph. Erdös and Gallai [4] first found necessary and sufficient conditions for a sequence of non-negative integers to be graphic and these conditions have since been refined by Hakimi [6] (stated in the sequel as Theorem 3.2) as well as (independently) by Havel [7]. Alternative characterisations and generalisations are due to Choudum [2], Sierksma & Hoogeveen [8] and Tripathi et al. [9], [10], [11]. This note states a result which is implicit in Hakimi [6], before giving an alternative characterisation of these necessary and sufficient conditions for a finite sequence of non-negative integers to be the degree sequence of a connected simple graph.

2. Preliminaries

Let $G = (V_G, E_G)$ be a graph where V_G denotes the vertex set of G and $E_G \subseteq [V_G]^2$ denotes the edge set of G (given that $[V_G]^2$ is the set of all 2-element subsets of V_G). An edge $\{a, b\}$ is denoted ab in the sequel. A graph is *finite* when $|V_G| < \infty$ and $|E_G| < \infty$, where $|X|$ denotes the cardinality of the set X. A graph is *simple* if it contain no loops (i.e. $a \neq b$ for all $ab \in E_G$) or parallel/multiple edges (i.e. E_G is not a multiset). The *degree* of a vertex v in a graph G, denoted $deg(v)$, is the number of edges in G which contain v. A *path* is a graph with n vertices in which two vertices, known as the *endpoints*, have

Email address: jonny_mclaughlin@hotmail.com (Jonathan McLaughlin)

degree 1 and $n-2$ vertices have degree 2. A graph is *connected* if there exists at least one path between every pair of vertices in the graph. A *tree* is a connected graph with n vertices and $n-1$ edges. K_n denotes the *complete graph* on n vertices. All basic graph theoretic definitions can be found in standard texts such as [1], [3] or [5]. All graphs in this note are undirected and finite.

3. Degree sequences and graphs

A finite sequence $s = \{s_1, ..., s_n\}$ of non-negative integers is called *realisable* if there exists a finite graph with vertex set $\{v_1, ..., v_n\}$ such that $deg(v_i) = s_i$ for all $i = 1, ..., n$. A sequence s which is realisable as a simple graph is called *graphic*. Given a graph G then the *degree sequence* of G, denoted $d(G)$, is the monotonic non-increasing sequence of degrees of the vertices in V_G. This means that every realisable (resp. graphic) sequence s is equal to the degree sequence $d(G)$ of some graph (resp. simple graph) G (subject to possible rearrangement of the terms in s). The maximum degree of a vertex in G is denoted Δ_G and the minimum degree of a vertex in G is denoted δ_G. In this note all sequences will have *positive* terms as the only connected graph which has a degree sequence containing a zero is $(\{v\}, \{\})$.

The following theorem states necessary and sufficient conditions for a sequence to be realisable (though not necessarily graphic).

Theorem 3.1 (Hakimi) *Given a sequence $s = \{s_1, ..., s_n\}$ of positive integers such that $s_i \geq s_{i+1}$ for $i = 1, ..., n-1$ then s is realisable if and only if $\sum_{i=1}^{n} s_i$ is even and $\sum_{i=2}^{n} s_i \geq s_1$.*

To address the issue of when a sequence is graphic, Hakimi describes in [6] a process he called a *reduction cycle* and uses it to state the following result.

Theorem 3.2 (Havel, Hakimi) *Given a sequence $\{s_1, ..., s_n\}$ of positive integers such that $s_i \geq s_{i+1}$ for $i = 1, ..., n-1$ then the sequence $\{s_1, ..., s_n\}$ is graphic if and only if the sequence $\{s_2 - 1, s_3 - 1, ..., s_{s_1+1} - 1, s_{s_1+2}, ..., s_n\}$ is graphic.*

4. Degree sequences and connected graphs

Definition 4.1 *A finite sequence $s = \{s_1, ..., s_n\}$ of positive integers is called connected (resp. connected and graphic) if s is realisable as a connected graph (resp. connected simple graph) with vertex set $\{v_1, ..., v_n\}$ such that v_i has degree s_i for all $i = 1, ..., n$.*

Of course disconnected realisations of connected and graphic degree sequences exist, for example, $(2,2,2,2,2,2)$ can be realised as a 6-cycle or as two disjoint 3-cycles.

As a graph is connected if and only if it contains a spanning tree, then a simple induction argument on the number of edges shows that every spanning tree of a graph G, with $|V_G| = n$, has exactly $n-1$ edges. Hence, a necessary condition for a graph G, with $|V_G| = n$, to be connected is that $|E_G| \geq n-1$.

The following theorem states necessary and sufficient conditions for a sequence to be connected but not necessarily simple.

Theorem 4.2 (Hakimi) *Given a sequence $s = \{s_1, ..., s_n\}$ of positive integers such that $s_i \geq s_{i+1}$ for $i = 1, ..., n-1$ then s is connected if and only if s is realisable and $\sum_{i=1}^{n} s_i \geq 2(n-1)$.*

The two main tools used in the proof of Theorem 4.2 are *d-invariant operations* (which leave degree sequences unchanged) and Lemma 4.3 (which appears in [6] as Lemma 1).

Consider a graph G with $d(G) = (d_1, ..., d_n)$. Given any two edges $ab, cd \in E_G$, where a, b, c and d are all distinct, then G is transformed by a *d-invariant operation* into G' when either

- $V_{G'} = V_G$ and $E_{G'} = (E_G \setminus \{ab, cd\}) \cup \{ac, bd\}$, or
- $V_{G'} = V_G$ and $E_{G'} = (E_G \setminus \{ab, cd\}) \cup \{ad, bc\}$.

Figure 1 shows both d-invariant operations.



Figure 1: The two possible d-invariant operations on G resulting in G' such that $d(G) = d(G')$

These d-invariant operations are used to prove the following important result.

Lemma 4.3 *Let $G_1, G_2, ..., G_r$, (with $r > 1$), be maximally connected subgraphs of G such that not all of the G_i are acyclic, then there exists a graph G' with $r-1$ maximally connected subgraphs such that $d(G') = d(G)$.*

The essence of Lemma 4.3 is presented in Figure 2. Note that *worst-case-scenarios* are assumed i.e. G_1 is a cycle and G_2 is acyclic with the graph $\big(V_{G_2}, (E_{G_2} \setminus \{cd\})\big)$ being disconnected.



Figure 2: G' has one less connected subgraph than G following a d-invariant operation

5. Results

The first result, Theorem 5.1, is an explicit statement of a result implicit in [6]. The second result, Theorem 5.2, is a new alternative characterisation of Theorem 5.1 and has a similar flavour to that of Theorem 3.2.

Theorem 5.1 *Given a sequence* $s = \{s_1, ..., s_n\}$ *of positive integers such that* $s_i \geq s_{i+1}$ *for* $i = 1, ..., n-1$ *then* s *is connected and graphic if and only if the sequence* $s' = \{s'_1, ..., s'_{n-1}\} = \{s_2 - 1, s_3 - 1, ..., s_{s_1+1} - 1, s_{s_1+2}, ..., s_n\}$ *is graphic and* $\sum_{i=1}^{n} s_i \geq 2(n-1)$.

Proof ($\Rightarrow$) Suppose that s is connected and graphic. It is required to show that s' is graphic and that $\sum_{i=1}^{n} s_i \geq 2(n-1)$.
As $s = d(G)$ for some simple (connected) graph G then $s_i \leq n-1$ for all $i = 1, ..., n$ and there exists a graph G' with vertex set $V_{G'} = V_G \setminus \{v_1\}$ and edge set $E_{G'} = E_G \setminus \{v_1 v_i \mid v_1 v_i \in E_G\}$ such that $d(G') = s'$. As G is a simple graph then it follows that G' is also a simple graph, hence s' is graphic. As $s = d(G)$ for some (simple) connected graph G, where $|V_G| = n$, then as G is connected $|E_G| \geq n-1$, hence $\sum_{i=1}^{n} s_i \geq 2(n-1)$.

($\Leftarrow$) Suppose that s' is graphic and that $\sum_{i=1}^{n} s_i \geq 2(n-1)$. It is required to show that s is both graphic and connected.

As s' is graphic then $s'_i \leq n-2$ for all $i \in \{1, ..., n-1\}$. Adding a term s_1 (which is necessarily $\leq n-1$) results in s also being graphic as in the worst case scenario i.e. where $s_1 = s_n = n-1$, then $s = \{s_1, ..., s_n\} = \{n-1, ..., n-1\}$ which is the degree sequence of the simple graph K_n. Suppose that $\sum_{i=1}^{n} s_i = 2(n-1)$, then s is the degree sequence of a graph G with $n-1$ edges and n vertices

which means that G is a tree, hence s is connected. If $\sum_{i=1}^{n} s_i > 2(n-1)$ then either $s = d(G)$ for some connected graph G or it is possible to apply Lemma 4.3 repeatedly until a graph G is found such that G is connected and $d(G) = s$. □

The following result is what can be thought of as a *connected version* of Theorem 3.2. However, note that it is not possible to simply add the word *connected* to the statement of Theorem 3.2 as $s = (2, 2, 1, 1)$ is connected but $s' = (1, 0, 1)$ is not connected.

Theorem 5.2 *Given a sequence $s = \{s_1, ..., s_n\}$ of positive integers such that $s_i \geq s_{i+1}$ for $i = 1, ..., n-1$ then s is connected and graphic if and only if the sequence $s' = \{s'_1, ..., s'_{n-1}\} = \{s_1 - 1, s_2 - 1, ..., s_{s_n} - 1, s_{s_n+1}, ..., s_{n-1}\}$ is connected and graphic.*

Proof ($\Rightarrow$) Suppose that $s = \{s_1, ..., s_n\}$ is connected. It is required to show that s' is both graphic and connected.

To show that s' is graphic it is required to show that $\sum_{i=1}^{n-1} s'_i$ is even and that all vertices have degree less than or equal to $n-2$. Observe that

$$\sum_{i=1}^{n-1} s'_i = \sum_{i=1}^{n-1} s_i - s_n = \sum_{i=1}^{n} s_i - 2s_n.$$

As s is graphic then $\sum_{i=1}^{n} s_i$ is even and so $\sum_{i=1}^{n} s_i - 2s_n$ is also even. As s is graphic then all vertices $v_i \in V_G$ with $i \in \{1, ..., n\}$ must satisfy $deg(v_i) \leq n-1$. All vertices with degree $n-1$ in G are necessarily connected to v_n whereas vertices with degree less than $n-1$ may or may not be connected to v_n. It follows that after deleting v_n and all edges containing v_n that the maximum degree which any vertex can have in any G' is $n-2$ (where $d(G') = s'$).

To show that s' is connected it is required to show that $\sum_{i=1}^{n-1} s'_i \geq 2(n-2)$ i.e. there exists a graph G' with $d(G') = s'$ and $|V_{G'}| = n-1$ such that $|E_{G'}| \geq n-2$. As s is graphic then $1 \leq s_i \leq n-1$.

Let $s_n = 1$: As $s_n = 1 = \delta_G$ then $s' = (s_1 - 1, s_2, ..., s_{n-1})$. Not all $s_i = 1$ except in the case where $s = (1, 1)$ resulting in $s' = (0)$ which is a connected degree sequence. As s is connected and $deg(v_n) = 1$ then v_n is a leaf of a connected graph G and so deleting v_n cannot result in a disconnected graph G', hence s' is connected when $s_n = 1$.

Let $s_n = k$ where $2 \leq k \leq n-1$: As $s_n = k = \delta_G$ then

$$s' = (s_1 - 1, s_2 - 1, ..., s_{s_n} - 1, s_{s_n+1}, ..., s_{n-1}).$$

Assuming the worst case scenario i.e. $\Delta_G = \delta_G = k$, then this gives

$$s' = (\underbrace{k-1, ..., k-1}_{k}, \underbrace{k, ..., k}_{n-k-1})$$

which means that

$$\sum_{i=1}^{n-1} s'_i \geq k(k-1) + k(n-k-1) = k(n-2) \geq 2(n-2)$$

whenever $2 \leq k \leq n-1$. Hence, s' is connected when $s_n = k$ where $2 \leq k \leq n-1$.

($\Leftarrow$) Suppose that $s' = \{s'_1, ..., s'_{n-1}\} = \{s_1 - 1, s_2 - 1, ..., s_{s_n} - 1, s_{s_n+1}, ..., s_{n-1}\}$ is connected. It is required to show that s is both graphic and connected.

As s' is connected then there exists some G' with $|V_{G'}| = n-1$ and $d(G') = s'$ where the degree of all vertices in $V_{G'}$ is less than or equal to $n-2$. As $V_G = V_{G'} \cup \{v_n\}$ (and all s_i are necessarily $\leq n-1$) then all vertices in V_G will have degree at most $n-1$. Observe that

$$\sum_{i=1}^{n} s_i = \sum_{i=1}^{n-1} s'_i + 2s_n.$$

As s' is graphic then $\sum_{i=1}^{n-1} s'_i$ is even and so $\sum_{i=1}^{n-1} s'_i + 2s_n$ is also even.

As s' is connected then there exists some G' with $d(G') = s'$ where $|E_{G'}| \geq n-2$ as $|V_{G'}| = n-1$. As $s_n \geq 1$ then this means that there is at least one edge in G which has v_n as an endpoint and some v_i with $i \in \{1, ..., n-1\}$ as the other endpoint.



Figure 3: If s' is connected and graphic then s is connected and graphic

This observation along with the fact that $|E_G| \geq |E_{G'}| + 1 > n-2$, where $|V_G| = n$, means that s is connected. □

Example 5.3 *An example, using Theorem 5.2, of what Hakimi would term a "set of successive reduction cycles" is shown in Figure 4.*



Figure 4: A set of successive reduction cycles

6. Comments

Theorem 3.2 is used in [6] to check algorithmically when a given sequence s is graphic. In a similar manner, Theorem 5.2 suggests an algorithm which can be used to determine if a given sequence s is connected and graphic.

References

[1] J. A. Bondy and U. S. R. Murty. *Graph theory*, volume 244 of *Graduate Texts in Mathematics*. Springer, New York, 2008.

[2] S. A. Choudum. A simple proof of the Erdös-Gallai theorem on graph sequences. *Bull. Austral. Math. Soc.*, 33(1):67–70, 1986.

[3] R. Diestel. *Graph theory*, volume 173 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 2000.

[4] P. Erdös and T. Gallai. Graphs with prescribed degrees of vertices. *Mat. Lapok*, 11:264–274 (in Hungarian), 1960.

[5] R. Gould. *Graph theory*. The Benjamin/Cummings Publishing Co. Inc., Menlo Park, CA, 1988.

[6] S. L. Hakimi. On realizability of a set of integers as degrees of the vertices of a linear graph. I. *J. Soc. Indust. Appl. Math.*, 10:496–506, 1962.

[7] V. Havel. A remark on the existence of finite graphs. *Časopis Pěst. Mat.*, 80:477–480 (in Czech), 1955.

[8] G Sierksma and H Hoogeveen. Seven criteria for integer sequences being graphic. *J. Graph Theory*, 15(2):223–231, 1991.

[9] A. Tripathi, S. Venugopalan, and D. B. West. A short constructive proof of the Erdös and Gallai characterization of graphic lists. *Discrete Math.*, 310(4):843 – 844, 2010.

[10] A. Tripathi and S. Vijay. A note on a theorem of Erdös and Gallai. *Discrete Math.*, 265:417 – 420, 2003.

[11] A. Tripathi and S. Vijay. A short proof of a theorem on degree sets of graphs. *Discrete Appl. Math.*, 155(5):670 – 671, 2007.